

LESCO®

4 Flowable Mancozeb BROAD SPECTRUM FUNGICIDE



Active Ingredients

MANCOZEB[®]:

A coordination product of zinc ion and manganese

ethylene bisdithiocarbamate 37.00%

In which the ingredients are:

Manganese⁺⁺ 7.40%

Zinc⁺⁺ 0.90%

Ethylene bisdithiocarbamate
ion (C₄H₆N₂S₄)⁻ 28.70%

INERT INGREDIENTS 63.00%

TOTAL

100.00%

¹Equivalent to 4 lb active ingredient per gallon

KEEP OUT OF REACH OF CHILDREN CAUTION

Notice: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal instructions. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

FIRST AID

IF INHALED:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact Chem-Trec at 1-800-424-9300 for emergency medical treatment information.

TM

UCC 128
(01) 0075807380230 (3)

#084032

NET CONTENTS:
2.5 gal (9.46 l)

LESCO is a registered trademark and the sweeping design is a trademark of LESCO Technologies, LLC.
Rainshield is a trademark of Dow AgroSciences LLC. 03-131

EPA Reg. No. 62719-396-10404

Manufactured For: LESCO, Inc. • 1301 East 9th Street • Cleveland, OH 44114-1849

EPA Est. 5905-GA-01

MADE IN FRANCE

900-009730/

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

May Cause Irritation Of Nose, Throat, Eyes And Skin. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category **A** on an EPA chemical-resistance category selection chart.

Applicators and other handlers (other than mixers and loaders) must wear:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Mixers and Loaders must wear:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks
- Protective eyewear
- Chemical-resistant apron when cleaning equipment, mixing or loading

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove PPE immediately after handling this product.
- Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. For terrestrial uses, do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or disposing of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of **24** hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Seed treatments and professional applications to lawn grasses, golf courses, industrial (office park), municipal and residential lawns are not within the scope of the Worker Protection Standard.

- Keep unprotected persons out of treated area until sprays have dried.

STORAGE AND DISPOSAL

Storage: Keep from freezing. Store in a cool, well-ventilated area, but not below 32°F. Do not allow to become overheated in storage. This may bring on chemical changes that will impair the fungicidal effectiveness of LESCO 4 Flowable Mancozeb. Keep container closed when not in use.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

Container Disposal: Plastic Containers- Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Fiber Drum (with liner)- Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities. If drum cannot be reused, dispose of in the same manner.

Bulk Containers- Drain thoroughly and return to specified destination for cleaning and reuse.

Steps to be Taken in Case Material is Released or Spilled: Dike and contain the spill. Transfer liquid and solid diking material to separate containers for recovery or disposal. Flush contaminated area with a large amount of water to a chemical or sanitary sewer containing a settling pit. Remove contaminated clothing and wash affected skin areas with soap and water. Wash clothing before reuse. Keep the solids out of the municipal sewers and open bodies of water. Refer to Precautionary Statements.

GENERAL USE INFORMATION

LESCO 4 Flowable Mancozeb is a broad-spectrum protectant fungicide recommended for outdoor or greenhouse grown crops. Optimum disease control is achieved when the fungicide is applied in a regularly scheduled preventative spray program. The addition of a surfactant will improve fungicide performance by providing a more uniform spray deposit, increased foliar redistribution, and improved fungicide retention during periods of wet weather.

USE RATE DETERMINATION

Carefully read, understand, and follow label use rates and restrictions.

Under low disease conditions, minimum label rates per application can be used while maximum label rates and the minimum interval may be used for severe or threatening disease conditions.

For proper application to turf, determine the square footage to be treated, divide the footage by 1000, and multiply by the recommended fungicide use rate per 1000 sq ft, and then determine the amount of water required to provide adequate coverage. When treating ornamentals, determine the recommended fungicide use rate and the spray gallonage required to provide thorough coverage. Careful calibration of spray equipment is recommended prior to use. Prepare only the amount of spray solution to treat the desired area.

If only a portion of the container's contents are to be used, thoroughly shake the container prior to measuring. When small quantities of spray solution are being prepared for use in hand or power sprayers, the following conversion table should be followed (rates are based on dilute thorough coverage sprays):

Recommended Use Rate per Acre or per 100 Gallons ¹	Fluid Ounces of LESCO 4 Flowable Mancozeb Required for:			
	10 gal	5 gal	2 gal	1 gal
0.8 qts	2.6	1.3	0.5	0.3
1.0 qt	3.2	1.6	0.7	0.3
1.2 qt	3.8	1.9	0.9	0.3
1.6 qt	5.1	2.6	1.0	0.5
2.0 qt	6.4	3.2	1.3	0.6
2.4 qt	8.0	4.0	1.6	0.8
3.2 qt	10.3	5.1	2.1	1.0
4.8 qt	15.4	7.7	3.1	1.6

1 cup = 8 fluid ounces or 237 milliliters
1 fluid ounce = 2 tablespoons or 30 milliliters
1 tablespoon = 3 teaspoons or 15 milliliters
¹Dilute thorough coverage sprays.

MIXING

Be sure the sprayer is clean and not contaminated with other materials prior to use. When using an agitated spray tank fill tank 1/2 to 3/4 full with clean water and start agitation. Be certain that the agitation system is working properly and creates a rolling rippling on the liquid surface. With the agitator running add the required amount of LESCO 4 Flowable Mancozeb to the tank. Continue filling tank with the remainder of the water. When using a hand sprayer, premix LESCO 4 Flowable Mancozeb as a slurry in a small container before adding to the spray tank. Slowly pour the appropriate amount of LESCO 4 Flowable Mancozeb into a small container containing an equal volume of water while mixing. Mix until the LESCO 4 Flowable Mancozeb is thoroughly wetted. Add additional water if necessary to make solution flowable. Add the contents of the slurry tank to a 1/2 filled sprayer, continue filling tank with remainder of water and mix well. Always add LESCO 4 Flowable Mancozeb into solution prior to adding any additional materials to the tank.

COMPATIBILITY

LESCO 4 Flowable Mancozeb is compatible with most commonly used fungicides, insecticides and growth regulators. When preparing tank mixes, user should consult spray compatibility charts or State Cooperative Extension Service Specialists prior to actual use.

Spray Adjuvants

The addition of agricultural surfactants to LESCO 4 Flowable Mancozeb sprays will improve initial spray deposits, fungicide redistribution and weatherability.

Place LESCO 4 Flowable Mancozeb into suspension prior to adding an adjuvant to the spray mixture. Read and carefully observe the precautionary statements and all other information appearing on both product labels prior to spray preparation.

APPLICATION

Ground

Thorough coverage foliar sprays generally result in optimum disease control. To achieve good coverage use proper spray pressure, gallonage per acre, nozzles (generally hollow cone), disc (generally D-5 to D-7), nozzle spacing, and tractor speed. Consult spray nozzle and accessory catalogues for specific information on proper equipment calibration.

Hand Sprayers

Thoroughly spray plant foliage until runoff.

CHEMIGATION USE DIRECTIONS

Sprinkler Irrigation

LESCO 4 Flowable Mancozeb must be applied on a regular protectant fungicide schedule, **not an irrigation schedule**. If irrigation cycles are less frequent than recommended LESCO 4 Flowable Mancozeb application intervals, ground or aerial applications must supplement chemigation applications to achieve adequate disease control.

Apply LESCO 4 Flowable Mancozeb only through sprinkler irrigation systems including center-pivot, lateral move, end tow, side (wheel) roll, traveler, solid set, or hand move irrigations systems. Do not apply product through any other type of irrigation system.

Lack of fungicidal effectiveness or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialist, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water system are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Before applying LESCO 4 Flowable Mancozeb through sprinkler irrigation equipment, the chemigation system must meet the following specifications:

- Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- Chemigation systems connected to public water systems must contain a functional reduced-pressure zone (RPZ), backflow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- Systems not connected to a public water supply must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located in the irrigation pipeline to prevent water source contamination from back flow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.

Center-pivot, Lateral Move, End Tow, and Traveler Irrigation Equipment:(use only with electric or oil hydraulic drive systems, which provide a uniform water distribution)

- Determine size of area to be treated.
- Determine the time required to apply no more than 1/4 inch water (6,750 gallons water per acre) over the area to be treated when the system and injection equipment are operated at normal pressures recommended by the equipment manufacturer. Run system at 80 to 95% of manufacturer's rated capacity.
- Using only water, determine the injection pump output when operated at normal line pressure.
- Determine the amount of LESCO 4 Flowable Mancozeb required for the treatment area.
- Add the required amount of LESCO 4 Flowable Mancozeb and sufficient water to meet the injection time requirements of the solution tank.
- Maintain constant solution tank agitation during the injection period.
- Stop injection equipment after treatment is completed. Continue to operate the system until LESCO 4 Flowable Mancozeb solution has cleared the sprinkler head.

Solid-set, Side (wheel) Roll, and Hand Move Irrigation Equipment:

- Determine acreage covered by sprinkler.
- Fill injector solution tank with water and adjust flow rate to use contents over a 10 to 30 minute interval.
- Determine the amount of LESCO 4 Flowable Mancozeb required for the treatment area.
- Add the required amount of LESCO 4 Flowable Mancozeb into the same quantity of water used to calibrate the injection equipment.
- Maintain constant solution tank agitation during the injection period.
- Operate system at normal pressures recommended by the manufacturer of the injection equipment and used for the time interval established during calibration.
- Inject LESCO 4 Flowable Mancozeb at the end of the irrigation cycle or as a separate application to maximize foliar fungicide retention.
- Stop injection equipment after treatment is completed. Continue to operate the system until LESCO 4 Flowable Mancozeb solution has cleared the last sprinkler head.

Disease Monitoring

LESCO 4 Flowable Mancozeb is a broad-spectrum, protectant fungicide. If not applied on a routine protectant spray schedule, turf and ornamental plants should be frequently observed for disease sign or symptoms. Fungicide application should be made, at the recommended label use rate and spray schedule, at the first sign of disease, report of disease in the area, or during environmental conditions favorable for disease development.

RESTRICTIONS

Users should carefully read, understand, and follow all use restrictions prior to using LESCO 4 Flowable Mancozeb.

TURF

For golf courses, sod farms, industrial or municipal turf areas and professional applications to residential lawns. **Not for use by homeowners.**

Start application when grass greens-up in spring or when disease first appears and repeat at 7 to 14 day intervals or until disease threat is past. When conditions are especially favorable for disease development, apply maximum fungicide use rate on a 7 day spray schedule. Apply in sufficient water to provide adequate coverage.

Turf Tolerance: Treated turfgrass should be maintained in a vigorous growing condition. Turfgrass under stress will not respond to fungicide treatments as well as well-maintained turfgrass. Turfgrass tolerance to this product has been found to be acceptable, however, this product and tank mixtures with other products have not been tested on all varieties of every turfgrass species or under all possible growing conditions. If user is unfamiliar with the performance of Lesco 4 Flowable Mancozeb or tank mixtures, under user growing conditions, a limited area of turfgrass should be treated prior to initiating large-scale applications. The user should always exercise reasonable judgment and caution when using this product.

Crop	Diseases Controlled	LESCO 4 Flowable Mancozeb Rate per Application (fl oz/1000 sq ft)	Remarks (Also Refer to Directions for Use)	Restrictions:
Assorted Grasses	Helminthosporium melting-out Rust (leaf, stem, stripe)	6.4		Do not graze treated areas. Do not use on grasses intended for grazing, such as range or pasture grasses. Do not feed clippings to livestock. Do not use on grasses grown for seed.
	Copper spot Fusarium blight Powdery mildew Red thread Slime mold	6.4 to 12.8		
	Algae	9.6		
	Dollar spot	9.6 to 12.8		
	Rhizoctonia brown patch	6.4	Apply on a 7 day spray schedule.	
	Pythium blight	12.8	Apply at 5 day intervals, or more frequently, if conditions are especially favorable for disease development	
	Fusarium snow mold	9.6 to 12.8	Apply at 2 to 6 week intervals during winter.	
	Gray leaf spot	12.8	Apply on a 14 day spray schedule when conditions are favorable for disease development.	

ORNAMENTALS

Not intended for use on fruit trees by homeowners. Neither the manufacturer nor the seller has determined the effects of using LESCO 4 Flowable Mancozeb on ornamentals not specified on this label. Prior to any large-scale applications on such plants, the user should determine the effects of LESCO 4 Flowable Mancozeb by testing a small section of the type of plants treated. User assumes all risk arising out of application to unlabeled plants. The Conditions of Sale and Warranty apply to all uses.

For outdoor or greenhouse use, apply the equivalent of 1.2 qt LESCO 4 Flowable Mancozeb per 100 gal dilute spray (1.2 quarts of LESCO 4 Flowable Mancozeb per acre).

Begin spraying when plants are well leaved out or at first sign of disease, in a full coverage spray at 7- to 10-day intervals throughout season or follow State Extension Service recommendations for disease control on the following ornamental plants.

Crop	Diseases Controlled	Remarks (Also Refer To Directions For Use)	Restrictions
African violet	Botrytis blight		Do not use for food or feed purposes (applies to all Ornamental uses).
Almond (Ornamental)	Leaf spot		
Alyssum	Leaf spot		
Anthurium	Anthraco-nose, spadix rot		
Apple (Ornamental)	Fabraea leaf spot Rust Scab		
Arborvitae	Cercospora blight		
Areca Palm	Leaf spot		
Ageratum	Botrytis blight, Rust		
Ash, mountain	Entomosporium leaf spot Guignardia leaf blotch		
Ash, white	Anthraco-nose Cylindrosporium leaf spot		
Aster	Leaf spot		
Aster, perennial	Puccinia rusts		
Aucuba, japonica	Alternaria leaf spot Anthraco-nose		
Azalea	Cylindrocladium rot Petal blight Phytophthora twig and bud blight	Apply in a full coverage spray, 2 to 3 times a week, while flowers are opening. Direct spray into flowers and thoroughly spray ground under bushes.	
Begonia	Botrytis blight		
Boxwood	Leaf spot		
Buffaloberry	Cylindrosporium leaf spot		
Camellias	Petal blight	Refer to Azalea.	

ORNAMENTALS (cont.)

Crop	Diseases Controlled	Remarks (Also Refer To Directions For Use)	Restrictions
Carnation	Rust Septoria leaf spot		
Cedar, red (Juniper)	Cercospora blight Phomopsis blight		
Chrysanthemum	Ascochyta blight Botrytis petal spot Rust	Apply twice weekly during blooming period.	
Cockscomb (Celosia)	Alternaria leaf spot		
Conifers(Christmas Trees)	Lophodermium needle cast Pine gall rust Scirrhia brown spot	Begin application in spring or early summer before infection occurs. Repeat after heavy rains and at two- week intervals as long as needed.	
Cordyline	Cercospora leaf spot		
Crabapple (Ornamental)	Cedar-apple rust Scab Sphaeropsis leaf spot		
Cypress, Arizona (Cupressus spp.)	Cercospora blight Monochaetia canker		
Dahlia	Botrytis blight		
Delphinium	Botrytis blight		
Dieffenbachia	Leptosphaeria brown spot		
Dogwood, flowering	Anthracnose Elsinoe leaf spot Septoria leaf spot	Apply when buds begin to open, when bracts have fallen, 4 weeks later and again in late summer after flower buds for next season have formed.	
Dracaena	Fusarium leaf spot		
Elm	Black leaf spot		
Euonymus	Anthracnose		
Fatsia	Anthracnose		
Fern	Rhizoctonia blight		
Ficus	Cercospora leaf spot		
Fig	Cylindrocladium leaf spot		
Firethorn (Pyracantha)	Fusicladium scab		
Fir, Douglas	Swiss needle cast		
Fir, Fraser	Swiss needle cast		
Fuchsia	Botrytis blight Rust		

ORNAMENTALS (cont.)

Crop	Diseases Controlled	Remarks (Also Refer To Directions For Use)	Restrictions
Geranium	Rust		
Gladiolus	Botrytis blossom blight Curvularia leaf spot	Make regular weekly applications starting before diseases appear and increase to 2 or 3 applications per week during periods of heavy disease and during rainy weather. On flower spikes, reduce spray concentration to 0.6 qt per 100 gallons.	
Gloxinia	Botrytis blight		
Gypsophila	Botrytis blight		
Hawthorn	Cedar-apple rust Fabraea leaf spot Frogeye leaf spot Hawthorn rust Scab		
Hickory	Gnomonia leaf spot		
Holly	Purple spot		
Hollyhock	Anthracnose Cercospora leaf spot Puccinia rust		
Honeysuckle	Herpobasidium blight		
Horsechestnut, Buckeye	Alternaria leaf spot Guignardia leaf blotch		
Hydrangea	Botrytis blight Cercospora leaf spot		
Impatiens	Botrytis blight		
Iris	Didymellina leaf spot Mycosphaerella leaf spot Mystrosporium ink spot	(formerly Didymellina)	
Juniper	Phomopsis blight		
Larkspur	Rust		
Laurel, mountain	Cercospora leaf spot Petal blight	Refer to azalea.	
Ligustrum	Cercospora leaf spot		
Lily	Botrytis blight		
Magnolia	Gloeosporium leaf spot		
Maple	Alternaria leaf spot Phyllosticta leaf spot		Do not use on Sugar maples intended for the production of maple syrup.

ORNAMENTALS (cont.)

Crop	Diseases Controlled	Remarks (Also Refer To Directions For Use)	Restrictions
Marigold	Botrytis blossom blight	Do not use on French dwarf double or Signet type marigold seedlings.	
Narcissus	Botrytis blight (fire) Smoulder		
Oak	Actinopelte leaf spot Taphrina leaf blister		
Orchid (Dendrobium)	Botrytis blossom blight		
Oxalis	Rust		
Pachysandra	Volutella blight	Use a drenching spray of 1.6 qt qt per 50 gallons of water per 5,000 sq ft of bed. Start applications at first sign of disease and apply at least 5 applications at 10- to 14-day intervals.	
Pansy	Anthracnose		
Pears (Ornamental)	Fabraea leaf spot Rust Scab		
Peony	Botrytis blossom blight Phytophthora blight	Apply in early spring and early fall, drenching soil around plants as well as the foliage. Promptly destroy all infected plant parts.	
Peperomia	Cercospora leaf spot		
Petunia	Botrytis blight		
Philodendron	Dactylaria leaf spot Phytophthora leaf spot		
Phlox	Leaf spot		
Photinia	Entomosporium leaf spot		
Pine, Australia	Cyclaneusma needle cast		
Pine, Scotch	Cyclaneusma needle cast Gall rust		
Pittosporum	Alternaria leaf spot		
Pleomele	Fusarium leaf spot		
Poinsettia	Sphaceloma scab		
Poplar	Rust		
Primrose	Botrytis blight		
Protea	Botrytis blight		
Quince (Ornamental)	Fabraea leaf spot Rust Scab		

ORNAMENTALS (cont.)

Crop	Diseases Controlled	Remarks (Also Refer To Directions For Use)	Restrictions
Rhododendron	Cercospora leaf spot Discosia leaf spot Petal blight	Refer to azalea.	
Rose	Black spot Cercospora leaf spot Rust		
Rosemary	Rhizoctonia Aerial blight		
Schefflera	Alternaria blight		
Scotts Pine	Needle cast		
Skunkbush, sumac	Cylindrosporium leaf spot		
Snapdragon	Rust		
Spathiphyllum	Myrothecium leaf spot		
Statice	Cercospora frog-eye		
Strawflower	Rust		
Syngonium	Cephalosporium leaf spot		
Thorn Apple	Rust		
Tulip	Botrytis blight (fire)		
Venus, flytrap	Anthracnose		
Viburnum	Downy mildew Ramularia leaf spot		
Walnut	Anthracnose		Do not use treated walnuts for food or feed purposes.
Zinnia	Alternaria leaf blight		

CONDITIONS OF SALE AND WARRANTY

LESCO, Inc. warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label only when used in accordance with label directions under normal conditions of use. **LESCO, Inc. MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR PARTICULAR USE.** Handling, storage, and use of product by Buyer or User are beyond the control of LESCO, Inc. and Seller. Risk such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pests, drift to other crops or property or failure to follow label directions will be assumed by Buyer or User. **IN NO CASE WILL LESCO, Inc. OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM HANDLING, STORAGE OR USE OF THIS PRODUCT.**

ATTENTION: This product contains mancozeb and ETU, chemicals known to the State of California to cause cancer. ETU is also known to the State of California to cause birth defects or other reproductive harm.

LESCO is a registered trademark and the sweeping design is a trademark of LESCO Technologies LLC. 03-131

Rainshield is a trademark of Dow AgroSciences LLC (050902)

Made in France

Manufactured For:

LESCO, Inc.

1301 East 9th Street

Cleveland, OH 44114-1849

Net Contents: 2.5 gal (9.46L)

#084032

LESCO®



4 Flowable Mancozeb BROAD SPECTRUM FUNGICIDE

Notice: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal instructions. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

PEEL FILM HERE

Active Ingredients

MANCOZEB¹:

A coordination product of zinc ion and manganese

ethylene bisdithiocarbamate 37.00%

In which the ingredients are:

Manganese²⁺ 7.40%

Zinc²⁺ 0.90%

Ethylene bisdithiocarbamate
ion (C₄H₆N₂S₄)²⁻ 28.70%

INERT INGREDIENTS 63.00%

TOTAL 100.00%

¹Equivalent to 4 lb active ingredient per gallon

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

May Cause Irritation Of Nose, Throat, Eyes And Skin. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers (other than mixers and loaders) must wear:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Mixers and Loaders must wear:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks
- Protective eyewear

- Chemical-resistant apron when cleaning equipment, mixing or loading

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove PPE immediately after handling this product.
- Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

IF INHALED:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for further treatment advice.

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact Chem-Trec at 1-800-424-9300 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. For terrestrial uses, do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or disposing of wastes.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

UCC 128
(01) 0075807380230 (3)

#084032

Refer to label booklet for Directions for Use including Storage and Disposal

NET CONTENTS: 2.5 gal (9.46 l)

TM

LESCO is a registered trademark and the sweeping design is a trademark of LESCO Technologies, LLC.
Rainshield is a trademark of Dow AgroSciences LLC. 03-131 (050902)

EPA Reg. No. 62719-396-10404

Manufactured For: LESCO, Inc. • 1301 East 9th Street • Cleveland, OH 44114-1849 EPA Est. 5905-GA-01

MADE IN FRANCE

900-009730/

Base
6¹/₄" x 6⁷/₈"; Image Area: 6" x 4⁷/₈"
Minimum text size: 6 pt.

Photo Comp Corporation
401 East Main Street
Brownsburg, IN 46112
1-800-358-8973
e-mail: das@photocomp.com