

Mauget® IMISOL™

Systemic Insecticide / Fungicide in Ready to Use Capsules
For Tree Injection Use For Seasonal Supression of Certain
Insects and Diseases of Ornamental Trees

ACTIVE INGREDIENT:

Debacarb
[2-(2-ethoxyethoxy)ethyl-2-benzimidazole carbamate]1.7%
Carbendazim
Methyl 2-benzimidazole carbamate..... 0.3%
Imidacloprid
1-[(6-Chloro-3-pyridinyl)methyl -N-nitro-2 imidazolidinimine...5.0
%
INERT INGREDIENTS:93.0%
100.0%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID	
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, perferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advise.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off containated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-535-5053 for emergency treatment information.	
NOTE TO PHYSICIAN	
There is no specific antidote available. Treat Patient symptomatically.	

Mfg. By: J.J. Mauget Co.
Town, State: Arcadia, CA 91006
EPA Reg. No.: 7946-21
EPA Est. No.: 7946-CA-1

Net Contents:

_____ 288 capsules @ 4mL each, 1152mL net; 288 feeder tubes
24 capsules plus 24 feeder tubes per carton
_____ 24 capsules @ 4 mL, 96 mL net or
_____ 24 capsules @ 6 mL, 144 mL net
Shipping box: 12 cartons as above.

Rev.1/19/04D

GENERAL DIRECTIONS

Measure the tree at chest height in inches. If measuring the circumference of the tree, divide this number by six (6) to determine the number of capsules needed. If measuring the diameter of the tree, divide this number by two (2) to determine the number of capsules needed. Apply the injector units around the tree at the root flare. Use the following dosage depending upon trunk diameter at breast height (DBH).

4 mL - 2 to 10 inches dbh

6 mL - 10 inches dbh and above

Consult the enclosed pamphlet "Directions for use and application of Mauget Micro-Injection System" for additional instructions. Applicators shall remove capsules promptly after treatment.

Important: Preventative application is more effective than therapeutic treatment in trees showing disease symptoms. Trees in advanced stages of disease development may not respond to treatment.

Healthy trees under optimum conditions will absorb the full contents promptly. Infected trees will absorb the material more slowly due to the vascular plugging caused by the disease. If Imisol is not absorbed within 24 hours, the tree is considered high risk and has a poor chance of survival.

RESTRICTIONS

Do not inject trees that are less than two inches in diameter. This product is not to be used on trees which will produce food within the year following treatment.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool dry place out of the reach of children. Do not store below 45° F. Store capsules in an upright position in closed container. Keep out of direct sunlight when possible.

DISPOSAL: Dispose of empty capsules in a sanitary landfill or by incineration if approved by State and Local authorities.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A

MANNER INCONSISTENT WITH ITS LABELING.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC
ANIMALS**

CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Causes eye irritation. Applicator must wear water proof gloves and face shield when applying this product. Wash thoroughly with soap and water after handling. Avoid breathing vapors. Remove contaminated clothing and wash before reuse.

Personal Protective Equipment:

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:


- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical resistant gloves, such as polyethylene or butyl rubber or neoprene rubber of viton
- Protective eyewear

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.



NOTICE OF WARRANTY

J.J. Mauget Co. makes no warranty of merchantability, fitness for any purpose or otherwise expressed or implied concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

IMISOL™

Disease Treatment: Make application when disease first appears. Repeat treatment if disease symptoms progress. Do not repeat within 3 months of first application. Some diseases may require repeated yearly application. Preventative Dutch Elm Disease treatments are made 4 weeks after Bark Beetle emerges. Therapeutic treatments are made as soon as possible after flagging branch is observed. **IMISOL** Insecticide/fungicide is for use on ornamental trees for the control of the following insects: Adelgids, Aphids, Black Vine Weevil Larva, Bronze Birch Borer, Cottonwood Borer, Elm Leaf Beetle, Eucalyptus Longhorned Borer, Flatheaded Borer (including Alder and Birch Borer and Emerald Ash Borer), Japanese Beetle, Lacebugs, Leafhoppers, Leafminers, Mealybugs, Pine Tip Moth Larvae, Psyllids (including Lerp Psyllid), Royal Palm Bugs, Scale Insects (including Asian Cycad Scale), Thrips, Whiteflies, Douglas Fir Gall Midge, Douglas Fir Cone Moth Larvae and the following diseases:

USE	DISEASE
Alder	Ceratocystis Canker, Fusarium Wilt , Physalospora (Bleeding Canker)
Arborvitae	Kabatina Branch Canker
Ash	Anthraco nose (Fall application only)
Bay Tree	Fusarium Wilt, Nectria Canker
Birch	Melanconium Dieback
Buckthorn	Nectria (Tubercularie) Canker
Camphor	Verticillium Wilt
Carob	Verticillium Wilt
Catalpa	Verticillium Wilt
Cedar	Coryneum Blight, Diplodia Tip Blight, Phomopsis (Kabatina)
Cypress	Cedar Branch Canker
Douglas Fir	Phomopsis Canker
Elm	Cephalosporium Ulmi (Elm Wilt), Ceratocystis Ulmi (Dutch Elm Disease), Cytospora Canker, Fusarium Wilt, Vermicularia Dieback, Verticillium Wilt
Fir	Cytospora Canker
Gum (Sweet)	Ceratocystis Canker
Madrone	Thielaviopsis Decline, Fusarium Wilt
Magnolia	Ceratocystis Canker
Maple	Verticillium Wilt
Mimosa	Fusarium Perniciosum (Mimosa Wilt)
Mulberry (Fruitless)	Ceratocystis Canker
Oak	Oak Wilt (Ceratocystis fagacearum), Anthracnose (Spring application only), Nectria Canker, Oak Decline, (BotryodiplodiaCephalosporium Dothiorella Fusarium, Fusarium Pestalotia Verticicladiella Verticillium)
Olive	Fusarium Wilt, Verticillium Wilt
Palm	Penicillium Vermoeseni (Pink Bud Rot), Atropellis Canker, Ceratocystis Dieback
Pine	Fusarium moniliforme F. Subglutinans (Pine Pitch Canker), Leptographium Canker
Pistachio(Non-Crop)	Verticillium Wilt
Poplar	Cytospora Canker
Redwood (Coast and Sierra)	Botryosphaeria Branch Canker, Coryneum Canker
Spruce	Cytospora Canker
Sycamore	Anthraco nose (Spring application only), Ceratocystis Canker
Walnut (Black Non-Crop)	Melanconium Dieback
Willow	Cytospora Canker