

The following compatibility test should always be performed prior to full-scale mixing.

1. Pour 18 ounces of water into a quart jar.
2. Add 1 ounce of either the liquid fertilizer or liquid iron to be used.
3. Add 1 ounce of LESCO Eliminate Liquid Selective Herbicide.
4. Close jar and shake well.
5. Watch the mixture for several seconds after shaking and check again after 30 minutes.
6. If the mixture does not show signs of separating, the combination may be used. If the mixture foams excessively, gas, separates or gets very thick, do not combine for field application.
7. Compatibility may be improved by the use of a compatibility agent. Some suggested compatibility agents to try are Katio Laboratories Complex, Hopkins Chemical's Unite, Farm Chemicals Inc.'s Compat, Harcross Chemicals T-Mulz 734-2, Rigco Company's Rigco Compatibility Agent, Witco Chemical's Sponto 168D, Amoco Oil's Amoco Spray Mate and Universal Co-op's Chem-Link. These agents are all used in the same manner. Follow the previously outlined test procedures and add 1/6 ounce of the compatibility agent between steps (the compatibility agent must be added to the fertilizer or iron before adding the LESCO Eliminate Liquid Selective Herbicide).
8. If the mixture does not separate, gel, foam or get very thick, it may be used for field application. Mix only the amount to be sprayed. Do not allow to stand overnight.

STORAGE AND DISPOSAL

STORAGE: Always store pesticides in a secured warehouse or storage building. Do not store near open containers of herbicides and other pesticides, fertilizer and seed. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well-ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

PRODUCT DISPOSAL: Pesticide wastes are acutely hazardous. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY

LESCO, Inc. warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixtures with other chemicals and factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use, including the use of this product on turf species not recommended on this label.

ESCO is a registered trademark of LESCO Technologies, LLC
 Eliminate is a trademark of LESCO Technologies, LLC
 Herbi is a trademark of North American Micron.

CTRLPRD#06250625589.LBLREV.02/05/02 WPG



ELIMINATE™ LIQUID SELECTIVE HERBICIDE

FOR SELECTIVE BROADLEAF WEED CONTROL IN ORNAMENTAL LAWNS AND TURF GRASSES

CONTAINS MCPA, TRICLOPYR AND DICAMBA

CONTROLS: Dandelion, Chickweed, Black medic, Knotweed, Plantain, Oxalis, Clover, Cocklebur, Thistle and many other species of broadleaf weeds; some of which are listed on this label.

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	48.99%
Triethylamine Salt of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid**	5.59%
Dimethylamine Salt of Dicamba (3,6-Dichloro-o-Anisic Acid)***	4.82%
INERT INGREDIENTS:	40.60%
TOTAL	100.00%
Isomer Specific AOC Method, Equivalent to:	
*2-Methyl-4-Chlorophenoxyacetic Acid	40.0%
**3,5,6-Trichloro-2-Pyridinyloxyacetic Acid	0.38 lb/gal
***3,6-Dichloro-o-Anisic Acid	4.0%, 0.38 lb/gal

For Use By Professional Turf Maintenance Personnel, Landscaping
 or Commercial Applicators Only


**KEEP OUT OF REACH OF CHILDREN
 DANGER - PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
**SEE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS
 AND STATEMENT OF PRACTICAL TREATMENT**

Distributed by:
LESCO, INC.
 15885 Sprague Road
 Strongsville, Ohio 44136-1799

NET CONTENTS: 2.5 GAL (9.5 L)

#0625589



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER - PELIGRO**

Corrosive. This concentrate causes irreversible eye damage. Harmful if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing. Wear face shield or goggles when mixing and loading this product. After product has been diluted in accordance with Directions for Use, eye protection is not required. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

- IF IN EYES:** Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.
- IF ON SKIN:** Wash with plenty of soap and water. Get medical attention.
- IF SWALLOWED:** Call a doctor or get medical attention. Do not induce vomiting. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.
- NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes. Most cases of ground water contamination involving phenoxy herbicides such as MCPA have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

LESCO Eliminate Liquid Selective Herbicide is for use on Lawns, Turf, Golf Courses (Fairways, Aprons, Tees* and Roughs), Parks, Cemeteries and similar non-crop areas. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production or for research purposes.

*Excluding Bentgrass Tees.

USE PRECAUTIONS

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Do not use on carpetgrass, Dichondra, nor on lawns or turf where desirable clovers are present. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from Spring or early Fall applications when weeds are actively growing. Use the higher rate for hard-to-control weeds. Do not exceed specified dosages for any area, be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 90°F.

Avoid application to Centipede or St. Augustine grasses unless turf injury can be tolerated. The suitable use of LESCO Eliminate Liquid Selective Herbicide on non-recommended turf species may be determined by treating a small area at any rate/acre which does not exceed 3 pints/acre. Prior to treatment of larger areas, the treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area. The user assumes the responsibility for any plant damage or other liability resulting from the use of this product on species not recommended on this label. For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) turf should not be mowed 1 to 2 days before and following application; (3) reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectively to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

WEEDS CONTROLLED BY LESCO ELIMINATE LIQUID SELECTIVE HERBICIDE

- Alfalfa, Amaranth, Aster, Beardsr, Bedstraw, Beggarick, Beggarweed, Bindweed, Bitter sneezeweed, Black medic, Bladder campion, Bracken fern, Broomweed, Buckhorn, Buckwheat, Buffalo, Burdock, Burcumber, Burdock, Bursage, Buttercup, Canada thistle, Carpetweed, Catnip, Catsear, Chamise, Chionodoxa, Chicory, Cinquefoil, Clover, Cocklebur, Coffeeweed, Cornflower, Cornspeedwell, Cotton, Daisy, Dandelion, Dock, Dog fennel, Dragonhead mint, Evening primrose, Field pennycress, Flabane, Florida pusley, Frenchweed, Galinsoga, Geranium, Goatsbeard, Goldenrod, Goosefoot, Gromwell, Ground Ivy, Halfshrub sundrop, Hawkweed, Headail, Heartleaf, drymary, Hemp sesbania, Hempenettle, Herbit, Horsepenitence, Horseweed, Ironweed, Jerusalem artichoke, Jimsonweed, Knapweed, Knapwood, Krawel, Knotweed, Kochia, Ladyshrub, Lambsquarters, Lespedeza, Little starwort, Mallow, Marc's tail, Marshelder, Matchweed, Mayweed, Milkweed, Morningglory, Mustard, Nightflowering catchfly, Nightshade, Oxalis (Stricia and corniculata), Parsley-piert, Pennywort, Peppergrass, Pigweed, Purslane, Purslane, Poison henlock, Poison Ivy, Poison oak, Pokeweed, Poorjoe, Prickly sida, Prickly lettuce, Puncturevine, Purslane, Raqweed, Red sorrel, Red clover, Redvine, Redstem, Rubberweed, Sheep sorrel, Shepherds-purse, Sicklepod, Smartweed, Snakeweed, Sowthistle, Speedwell, Spiderwort, Spikeweed, Spiny Spurge, Spurry, Spurweed, Starbur, Stinging nettle, Stitchwort, Stunkweed, Sunflower, Tansy, ragwort, Tassel, Teaweed, Texas blueweed, Thistle, Toadflax, Trumpet creeper, Velvetleaf, Veronica, Vetch, Waterhenlock, Waterhemp, Whitebush, Wild violet, Wild radish, Wild onion, Wild carrot, Wild lettuce, Wild garlic, Wild geranium, Wood sorrel, Wormwood, Yankeeweed, Yarrow, Yellow starthistle, Yellow rocket, and many other broadleaf weeds.

ORNAMENTAL LAWNS AND TURF

Apply LESCO Eliminate Liquid Selective Herbicide at the rate of 2 to 3 pints in 20 to 240 gallons of water per acre (0.73 to 1.10 fluid ounces in 0.5 to 5.5 gallons of water per 1,000 square feet) to control weeds growing in turf planted to Bluegrass, Fescue, Rye, Bent (excluding golf course greens and tees), Bahia, Bermuda and Zoysia.

Herbi™ Controlled Droplet Applicator: Add 1-1/2 to 2-1/4 pints of LESCO Eliminate Liquid Selective Herbicide to the 5 pint Herbi bottle, then fill with water to make 5 pints of mixture or substitute 1/2 pint of a surfactant for water while agitating the solution.

While walking at approximately 1 pace (3 feet) per second, spray entire contents over 33,000 square feet (3/4 of an acre). Do not overlap (double coverage) at edge of spray patterns. Reduced rates (use 1/2 of rate shown above) of LESCO Eliminate Liquid Selective Herbicide must be applied when grass is stressed from heat, drought, etc.

Controlled Droplet Applicators - (CDA), Atomizers, and Spinning Disk Applicators: Use LESCO Eliminate Liquid Selective Herbicide at the rate of 2 to 3 pints per acre (0.73 to 1.10 fluid ounces per 1,000 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment).

NOTE: For all grasses (1) Do not overlap spray patterns; (2) Use reduced rates if grass is stressed from heat, drought, etc.; and (3) Follow CDA equipment spray instructions.

COMPATIBILITY

LESCO Eliminate Liquid Selective Herbicide can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizers and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test (given below) is recommended prior to mixing in the application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.