

Granular Turf Fungicide

Contains 1.00% Bayleton® Fungicide. For Systemic Control of Certain Diseases on Turf

NOT FOR USE on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

SYSTEMIC FUNGICIDE GRANULES

FOR CONTROL OF LAWN DISEASE ON BLUEGRASS, FESCUE, RYEGRASS, BERMUDA GRASS, ZOYSIAGRASS, ST. AUGUSTINE GRASS AND BENTGRASS

FOR THE PREVENTION OR CURE OF DOLLAR SPOT, COPPER SPOT, POWDERY MILDEW, RED THREAD, RUSTS, BROWN PATCH, ANTHRACNOSE, POA ANNUA DECLINE, STRIPED SMUT, FUSARIUM BLIGHT, GRAY SNOW MOLD, PINK SNOW MOLD, FUSARIUM PATCH AND SOUTHERN BLIGHT.

ACTIVE INGREDIENT:

1-(4-chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)	
-2-butanone	1.00%
OTHER INGREDIENTS	99.00%
TOTAL	100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact Chem-Trec at 1-800-424-9300 for emergency medical treatment.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapor, or dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Wash contaminated clothing with soap and hot water before reuse.

ENVIRONMENTAL HAZARDS

Do not contaminate water when disposing of equipment washwaters. Do not use on crops grown for food or forage. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Apply this product only as specified on this label. Do not apply when weather conditions favor drift from target area. Do not use clippings or graze for animal feed.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store product in original bag in a cool, dry, locked place out of reach of children.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

APPLICATION

This product is absorbed rapidly and works systemically from within the plant. Rainfall or sprinkler irrigation soon after application will assure rapid systemic uptake. Control will be less effective on turf suffering from drought stress. Therefore in order to achieve maximum control, turf should be maintained in a vigorously growing state through good cultural practices. Apply when the turf is fully established and actively growing. Apply Preventative Rates prior to appearance of disease and/or when damage from disease is anticipated. Apply Curative Rate when disease is first observed.

NOTE: Apply the specified amount of LESCO Granular Turf Fungicide with a standard granular applicator calibrated to uniformly apply recommended amount of product per 1,000 sq ft. Make application when turf is dry and following the mowing operation. Do not mow until a minimum of 1/4 inch of water has been applied to the treated area.

WARRANTY

LESCO, Inc. warrants that this product conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, it is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of use.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept., 1301 East 9th Street, Suite 1300, Cleveland, Ohio 44114-1849, referring to the item number found on this bag.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-a1.com>.

LESCO is a registered trademark and the sweeping design is a trademark of LESCO Technologies, LLC. Bayleton is a registered trademark of Bayer Corporation. SCOTTS is a registered trademark of The Scott Company.

G:\REGUL\WPERIEVIEW\Aprvd Lbl - Word\Control Products\Bayleton\069292.doc
Rev.: 10/20/04 BM

#069292

F1560
Distributed by: **LESCO, Inc.**
1301 East 9th Street
Cleveland, OH 44114-1849

Recommended Applications – Turfgrass Disease Control

Disease	How much to use per 1,000 sq ft		Remarks
	Preventative	Curative	
Dollar Spot	1.5 lb	-	Preventative Rate: Apply recommended rate on 30-day intervals. Protective activity of product may extend for as long as 60 days dependent upon environmental conditions.
	0.75 lb	-	Preventative Rate: Apply recommended rate on 14-day intervals. Protective activity of product may extend for as long as 30 days dependent upon environmental conditions.
	-	3 lb	Curative Rate: To control existing infections, apply the curative rate. Subsequent applications should be applied on a preventative schedule and rate.
Copper Spot Powdery Mildew Red Thread (<i>Laetisara fuciformis</i>)			Preventative Rate: Apply at 15 to 30-day intervals. When environmental conditions favor light to moderate disease development, use a longer interval. Protective activity of this product can be greater than 30 days depending on environmental conditions.
Rusts Brown Patch (<i>Rhizoctonia solani</i>) (suppression)	1.5 lb	3 lb	Curative Rate: To control existing infections, apply the curative rate. Subsequent applications should be applied on a preventative schedule and rate.
Anthracnose	3 lb	-	Preventative Rate: Apply at 30-day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 45 days.
Poa annua Decline	1.5 lb	-	Preventative Rate: Apply at 30-day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 30 days.
	-	3 lb	Curative Rate: To control existing infections, apply the curative rate. Subsequent applications should be applied on a preventative schedule and rate.
Stripe Smut	Preventative Rate 3 lb		Make single application in Spring as grass growth begins.
Fusarium Blight (<i>Fusarium culmorum</i>) (<i>Fusarium poae</i>)	(Except California) 3 lb (California Only) 3 to 6 lb		Apply first application in mid-June or 30 days prior to time this blight normally becomes evident. Apply at 30-day intervals. For California Only: Apply at 3 lb rate on 15 day intervals or at the 6 lb rate on 30 day intervals.
Summer Patch (<i>Phialophora graminicola</i>)			Apply in fall 30 days prior to turf dormancy. If turf breaks dormancy during months a second application should be made. Do not apply over snow cover or when turf is dormant.
Gray Snow Mold (<i>Typhula incarnate</i>) (Except California)	6 lb		Apply before conditions favorable for infection occur. Repeat as needed at 60 to 90-day intervals. Do not apply over snow cover or when turf is dormant. Use high rate in turf with a history of severe yearly disease damage.
Pink Snow Mold Fusarium Patch (<i>Fusarium nivale</i>) Southern Blight (<i>Sclerotium rolfsii</i>) (Except California)	3 to 6 lb		Apply in late Fall before snow cover. Do not apply over snow. Use the high rate in areas with a history of severe snow mold damage.
Gray Snow Mold Pink Snow Mold (California Only)	Single Application 7.5 to 12 lb per 1,000 sq ft		

SUGGESTED SPREADER SETTINGS*

Spreader	Recommended Rates in lb per 1,000 sq ft					
	.75 lb	1.5 lb	3.0 lb	6.0 lb	7.5 lb**	12 lb**
LESCO Calibration Gauge	---	#11	#14	#17	---	---
SCOTTS® R8-A	---	H ½	J	K ½	---	---

*IMPORTANT: Spreader settings are only approximate. Age and condition of spreader may cause wide variations. Be sure to calibrate your spreader with each application. Refer to Spreader Manufacturer's Operational Manual.

** Note: Single application only, California only.

NET WEIGHT: 25 lb (11.3 kg)

EPA Reg. No. 10404-58

EPA Est. No. 82757-FL-001(S), 82757-OH-001(M)

First letter of lot code indicates manufacturing site.

