

# LESCO<sup>®</sup> MOMENTUM<sup>®</sup> FX<sup>2</sup> HERBICIDE

FOR THE CONTROL OF ANNUAL AND PERENNIAL BROADLEAF WEEDS  
IN ORNAMENTAL TURF.

CONTAINS 2,4-D, TRICLOPYR, AND FLUROXYPYR.

**ACTIVE INGREDIENTS:**

Triisopropanolamine Salt of 2,4-Dichlorophenoxyacetic Acid*	44.20%
Triethylamine Salt of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid**	3.86%
1-Methylheptyl Ester of Fluroxypyr: [(4-amino-3-5-dichloro-6-fluoro-2-pyridinyl)oxy]acetic Acid, 1-methylheptyl Ester***	4.20%

**OTHER INGREDIENTS:** ..... 47.74%

**TOTAL** ..... 100.00%

**By Isomer Specific AOAC Method, Equivalent to:**

*2,4-Dichlorophenoxyacetic Acid	23.70%, 2.254 lbs./gal.
**3,5,6-Trichloro-2-Pyridinyloxyacetic Acid	2.77%, 0.263 lbs./gal.
***[(4-amino-3-5-dichloro-6-fluoro-2-pyridinyl)oxy]acetic Acid.	2.92%, 0.278 lbs./gal.

**INTENDED FOR USE BY PROFESSIONAL TURF MAINTENANCE PERSONNEL,  
LANDSCAPING OR COMMERCIAL APPLICATORS ONLY.  
DO NOT SELL, DISTRIBUTE OR USE THE PRODUCT IN NASSAU AND SUFFOLK COUNTY IN NEW YORK.**

**KEEP OUT OF REACH OF CHILDREN  
DANGER - PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand this label, find someone to explain it to you in detail.)

**SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL  
PRECAUTIONARY STATEMENTS**

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300  
For Medical Emergencies Only, Call (877) 325-1840

**#088525**  
**NET CONTENTS: 2.5 gal (9.5 L)**

TM

EPA REG. NO. 228-447-10404

EPA EST. NO. 228-IL-1

Distributed by: **LESCO, Inc.** • 1301 East 9th Street • Cleveland, OH 44114-1849

LESCO is a registered trademark and the sweeping design is a trademark of LESCO Technologies, LLC.  
Momentum is a registered trademark and used by LESCO, Inc. with permission of the trademark owner.



**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
DANGER – PELIGRO**

Corrosive. Causes irreversible eye damage. Harmful if swallowed. Do not get in eyes or on clothing. If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical-resistant apron. If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

**FIRST AID**

<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>HOT LINE NUMBER</b>	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact CHEMTREC for transportation and spills (800) 424-9300; for Human health (800) 832-HELP (800-832-4357); and ASPCA for animal health (800) 345-4735.</p>	

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

**NON-WPS TURF USES:** Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) (in general only agricultural plant uses are covered by the WPS) must wear face shield, goggles, or safety glasses and long pants, long-sleeved shirt, socks, shoes and chemical-resistant gloves when mixing, loading or applying this product. It is recommended that safety glasses include front, brow, and temple protection. In addition to the clothing and eye protection listed above, commercial mixer/loaders must wear chemical-resistant in place of rubber gloves except when the product is to be applied to a golf course. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

**NON-WPS INDUSTRIAL USES:** When mixing, loading or applying this product or repairing or cleaning equipment used with this product, wear face shield, goggles, or safety glasses and chemical-resistant gloves, long-sleeved shirt, long pants, socks and shoes. It is recommended that safety glasses include front, brow, and temple protection. For aerial applicators in an enclosed cockpit and applicators applying this product from a tractor that has a completely enclosed cab, eye protection is not required. Wash hands, face and arms with soap and water as soon as possible after mixing, loading or applying this product. After work, remove all clothing and shower using soap and water. Do not reuse clothing worn during the previous day's mixing and loading or application of this product without cleaning first. Clothing must be kept and washed separately from other household laundry. Remove saturated clothing as soon as possible and shower.

**WPS USES:** Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) (in general, agricultural plant uses are covered) must wear: long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material, shoes plus socks, and protective eyewear, when cleaning equipment, mixing or loading. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned. If this container is over one gallon and less than five gallons, mixers and loaders who do not use a mechanical system (such as a probe and pump or spigot) to transfer contents of this container must wear coveralls or a chemical-resistant apron in addition to the other required PPE.

**Engineering Controls Statements:** If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### USER SAFETY RECOMMENDATIONS

##### Users Should:

- Wash hands, face and arms with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product.
- Wash the outside of gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Drift or runoff from treated areas may be hazardous to aquatic organisms and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target area. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other spraying. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these herbicides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

This pesticide should be used strictly in accordance with the drift and run-off precautions on this label in order to minimize off-site exposure.

Under some conditions this product may have a potential to run-off to surface water or adjacent land. Where possible, use methods which reduce soil erosion, such as no till, limited till and contour plowing; these methods also reduce pesticide run-off. Where feasible, use application techniques such as T-banding and in-furrow techniques which incorporate the pesticide into the soil. Use of vegetation filter strips along rivers, creeks, streams, wetlands, etc. or on the downhill side of fields where run-off could occur will minimize water run-off is recommended.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift. Avoid spraying during conditions of low humidity and/or high temperatures.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, chemical-resistant gloves made of any waterproof material, shoes plus socks and protective eyewear.

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

For turf use, the maximum number of broadcast applications per treatment site is 2 per year.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

This product is a broad-spectrum herbicide for control of actively growing annual and perennial broadleaf weeds. It is for use on established Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields, Roadsides, Golf Courses (Fairways, Aprons and Roughs) and similar turf areas; also for use on Sod Farms. Do not use on Golf Greens or Tees.

This product may be applied to the following established turfgrass species:

**Established Cool Season Turfgrasses**

<b>Common Name</b>	<b>Scientific Name</b>
Bentgrass †	<i>Agrostis species</i>
Bluegrass, Kentucky	<i>Poa pratensis</i>
Fescue, chewing	<i>Festuca rubra</i> var. <i>commutata</i>
Fescue, creeping red	<i>Festuca rubra</i>
Fescue, sheeps	<i>Festuca ovina</i>
Fescue, tall	<i>Festuca arundinacea</i>
Ryegrass, perennial	<i>Lolium perenne</i>

**Established Warm Season Turfgrasses††**

<b>Common Name</b>	<b>Scientific Name</b>
Bahiagrass †	<i>Paspalum notatum</i>
Bermudagrass †	<i>Cynodon dactylon</i>
Buffalograss	<i>Buchloe dactyloides</i>
Centipedegrass †	<i>Eremochloa ophiuroides</i>
Zoysiagrass †	<i>Zoysia japonica</i>
Zoysiagrass †	<i>Zoysia tenuifolia</i>
Fescue, tall (growing in warm season areas)	<i>Festuca arundinacea</i>

† Do not apply to these turfgrass species unless potential turfgrass injury can be tolerated. Also, when treating these turfgrass species, do not apply more than 3 pints per acre. To minimize grass injury, additional applications should be made at least four weeks apart. Avoid swath overlaps.

†† Do not treat warm season turfgrass with this product when mowing height is less than 1/2 inch.

**Note:** This product may discolor and/or stunt turf that is not well-established or is stressed or weakened due to unfavorable climatic conditions, temperature extremes, drought, nematodes, or other factors which damage or weaken turf. Apply this product only to healthy, well-established turf that has a well-anchored root system. Do not apply this product to grass species not listed on this label unless potential turf injury can be tolerated.

#### USE PRECAUTIONS

Do not apply this product directly to, or otherwise permit drift or spray mist to come into contact with cotton, citrus, grapes, tobacco, vegetables, flowers, ornamental plants, shrubs, trees, and other desirable plants. Do not pour spray solutions near desirable plants. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Spray drift can be reduced by spraying when wind velocity is low, keeping spray boom as low as possible, using nozzles that deliver larger spray droplets and minimize the production of fine spray particles, and by using higher spray volumes and lower pressures. Applications should be made only when hazards from spray drift are at a minimum. Very small quantities of spray, which may not be visible, may seriously injure susceptible plants. Do not spray when wind is blowing toward susceptible crops or ornamental plants near enough to be injured. Do not apply to exposed roots of shallow rooted trees and shrubs. Maximum kill of weeds will be obtained from Spring or early Fall applications when weeds are actively growing. Do not apply more than 4 pints per acre per application. Mow newly seeded turf at least twice before treatment with this product.

For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) turf should not be mowed 1 to 2 days before and following application; (3) reseed no sooner than 3 to 4 weeks after application of this product.

#### WEEDS CONTROLLED

This product is designed to control many broadleaf weeds such as:

Alder	Bull nettle	Curly indigo	Healall	Mugwort
Annual yellow sweet clover	Bull thistle	Dandelion	Heartleaf drymary	Morningglory
Artichoke	Burdock	Dead nettle	Hedge bindweed	Mouseear chickweed
Aster	Burning nettle	Dock	Hedge mustard	Musk thistle
Austrian fieldcress	Bur ragweed	Dogbane	Hemp	Mustard
Bedstraw	Burweed	Dogfennel	Henbit	Narrowleaf plantain
Beggartick	Buttercup	Elderberry	Hoary cress	Narrowleaf vetch
Biden	Canada thistle	English daisy	Hoary plantain	Nettle
Bindweed	Carolina geranium	Fall dandelion	Hoary vervain	Nutgrass
Bird vetch	Carpetweed	False dandelion	Honeysuckle	Orange hawkweed
Bitterweed	Catchweed bedstraw	False flax	Hop clover	Oxalis
Bitter wintercress	Catsear	False sunflower	Horsenettle	Oxeye daisy
Black-eyed Susan	Catnip	Fiddleneck	Horsetail	Parsley-piert
Black medic	Chickweed	Field bindweed	Indian mallow	Parsnip
Black mustard	Chicory	Field pansy	Ironweed	Pearlwort
Blackseed plantain	Cinquefoil	Flea bane (daisy)	Jewelweed	Pennycress
Blessed thistle	Clover	Flixweed	Jimsonweed	Pennywort
Blue lettuce	Cockle	Florida betony	Kochia	Peppergrass
Blue vervain	Cocklebur	Florida pusley	Knawel	Pepperweed
Box elder	Coffeebean	Frenchweed	Knotweed	Pigweed
Bracted plantain	Coffeeweed	Galinsoga	Lambsquarter	Pineywoods bedstraw
Brassbuttons	Common chickweed	Garlic mustard	Lespedeza	Plains coreopsis
Bristly oxtongue	Common mullein	Goathead	Locoweed	Plantain
Broadleaf dock	Common sowthistle	Goatsbeard	Lupine	Poison hemlock
Broadleaf plantain	Corn Chamomile	Goldenrod	Mallow	Poison ivy
Broomweed	Creeping jenny	Ground ivy	Marshelder	Poison oak
Buckhorn	Crimson clover	Gumweed	Matchweed	Pokeweed
Buckhorn plantain	Croton	Hairy bittercress	Mexicanweed	
Bulbous buttercup	Cudweed	Hairy fleabane	Milk vetch	
	Curly dock	Hawkweed	Milkweed bloodflower	

(continued)

**WEEDS CONTROLLED** (continued)

Poorjoe	St. Johnswort	Spurweed	Venice mallow	Wild parsnip
Povertyweed	Scarlet pimpernel	Stinging nettle	Veronica	Wild radish
Prickly lettuce	Scotch thistle	Stinkweed	Vervain	Wild rape
Prickly sida	Sheep sorrel	Stitchwort	Vetch	Wild strawberry
Primrose	Shepherds-purse	Strawberry clover	Virginia buttonweed	Wild sweet potato
Prostrate knotweed	Slender plantain	Sumac	Virginia creeper	Wild vetch
Prostrate pigweed	Smallflower galinsoga	Sunflower	Virginia pepperweed	Willow
Prostrate spurge	Smartweed	Sweet clover	Wayleaf bullthistle	Witchweed
Prostrate vervain	Smooth dock	Tall nettle	Western clematis	Woolly morningglory
Puncture vine	Smooth pigweed	Tall vervain	Western salsify	Woodsorrel
Purslane	Sneezeweed	Tansy mustard	White clover	Woolly croton
Ragweed	Southern wild rose	Tansy ragwort	White mustard	Woolly plantain
Red clover	Sowthistle	Tanweed	Wild mustard	Wormseed
Redroot pigweed	Spanishneedle	Tarweed	Wild aster	Yarrow
Red sorrel	Spatterdock	Thistle	Wild buckwheat	Yellow rocket
Redstem filaree	Speedwell	Tick trefoil	Wild carrot	Yellowflower
Rough cinquefoil	Spiny Amaranth	Toadflax	Wild four-o'clock	pepperweed and
Rough fleabane	Spiny cocklebur	Trailing Crownvetch	Wild garlic	other broadleaf
Roundleafed marigold	Spotted catsear	Tumble mustard	Wild geranium	weeds.
Rush	Spotted knapweed	Tumble pigweed	Wild lettuce	
Russian pigweed	Spotted spurge	Tumbleweed	Wild marigold	
Russian thistle	Spurge	Velvet leaf	Wild onion	

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. Some hard-to-control broadleaf weeds may require a second broadcast treatment in 3 to 4 weeks. In the event more than two broadcast applications per site per year are necessary, use as a spot treatment. To spot treat, make up solution as listed below.

**RECOMMENDED MIXING INSTRUCTIONS**

Fill spray tank with water. Start agitation and slowly add the recommended amount of concentrate. Maintain continuous agitation after mixing and during application. If this product is allowed to stand in tank for extended periods of time, some separation may occur. Reagitate before use.

**TURF APPLICATION DIRECTIONS**

Apply 3 to 4 pints of this product per acre (1.1 to 1.5 oz./1,000 sq.ft.) to control above listed broadleaf weeds.

The maximum use rate for most warm season grasses is 3 pints per acre. For exceptions, see listing of warm season grasses listed above.

Make application using equipment that will ensure uniform coverage (see specific application directions below). Foliar sprays should be applied during warm weather when weeds are actively growing. Application under drought conditions may provide less than desirable results. When hard-to-control species are prevalent or when weeds are mature, use the higher rates.

On closely mowed golf course fairway bentgrass, apply this product at a maximum rate of 2 pints in 20 to 240 gallons of water per acre (0.73 fluid ounces in 0.5 to 5.5 gallons of water per 1,000 square feet). During conditions which promote turf stress use lower rates, 1 to 1-1/2 pints in 20 to 240 gallons of water, per acre (0.36 to 0.55 fluid ounces in 0.5 to 5.5 gallons per 1,000 square feet). Slight turf yellowing will disappear after one week. To minimize grass injury, a second application should not be made for at least 4 weeks. Avoid swath overlaps. Do not use on golf course greens or tees.

**Note:** Broadleaf weeds germinate at different times of the year. Only emerged weeds present at the time of application will be controlled by this product.

**Standard Volume Broadcast Applications:** Apply 3 to 4 pints of this product in enough water to deliver 20 to 200 gallons of total spray mix per acre (0.5 to 5 gallons spray per 1,000 square feet). Higher application volumes may be used when this product is tank mixed with liquid fertilizers.

**Low Volume Broadcast Applications:** Apply 3 to 4 pints of this product in enough water to deliver from 5 to 40 gallons of total spray mixture per acre. Use low pressures and application equipment capable of delivering a uniform droplet size that can wet the weed leaf surface.

**Targeted Zone Applications or Spot Treatment Using Portable Sprayers:** Mix 1.1 to 1.5 ounces of this product in enough water to make one gallon of spray and apply at any time broadleaf weeds are actively growing. Target plants should be sprayed to wet but not beyond point of runoff.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Always store pesticides in a secured warehouse or storage building. Do not store near open containers of herbicides and other pesticides, fertilizer and seed. Protect product from freezing. Store at temperatures above 25°F. If allowed to freeze, remix before using. This does not alter the product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

**PRODUCT DISPOSAL:** Pesticide wastes are acutely hazardous. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL FOR NON-REFILLABLE CONTAINERS:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY:**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Desirable plant injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of LESCO. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold LESCO harmless for any claims relating to such factors.

LESCO warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use.

**LESCO MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT.** Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) LESCO, and buyer assumes the risk of any such use.

To the extent allowed by law, LESCO shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF LESCO FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR AT THE ELECTION OF LESCO, THE REPLACEMENT OF PRODUCT.

This condition of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

LESCO is a registered trademark and the sweeping design is a trademark of LESCO Technologies, LLC.  
Momentum is a registered trademark and used by LESCO, Inc. with permission of the trademark owner.

(RV030206)

5/17/05 LSL  
1/26/06 LAL



# LESCO Momentum® FX<sup>2</sup> Herbicide

FOR THE CONTROL OF ANNUAL AND PERENNIAL BROADLEAF WEEDS IN ORNAMENTAL TURF.

CONTAINS 2,4-D, TRICLOPYR, AND FLUROXYPYR.

INTENDED FOR USE BY PROFESSIONAL TURF MAINTENANCE PERSONNEL, LANDSCAPING OR COMMERCIAL APPLICATORS ONLY.

**ACTIVE INGREDIENTS:**

Trisopropanolamine Salt of 2,4-Dichlorophenoxyacetic Acid* . . . . .	44.20%
Triethylamine Salt of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid** . . . . .	3.86%
1-Methylheptyl Ester of Fluroxypyr: [(4-amino-3-5-dichloro-6-fluoro-2-pyridinyl)oxy]acetic Acid, 1-methylheptyl Ester*** . . . . .	4.20%

**OTHER INGREDIENTS:**

.....	47.74%
<b>TOTAL:</b> .....	100.00%

**By Isomer Specific AOAC Method, Equivalent to:**

*2,4-Dichlorophenoxyacetic Acid. . . . .	23.70%, 2.254 lbs/gal
**3,5,6-Trichloro-2-Pyridinyloxyacetic Acid . . . . .	2.77%, 0.263 lbs/gal
***[(4-amino-3-5-dichloro-6-fluoro-2-pyridinyl)oxy]acetic Acid . . . . .	2.92%, 0.278 lbs/gal

**DO NOT SELL, DISTRIBUTE OR USE THE PRODUCT IN NASSAU AND SUFFOLK COUNTY IN NEW YORK.**

## KEEP OUT OF REACH OF CHILDREN DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand this label, find someone to explain it to you in detail.)  
**SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
DANGER - PELIGRO**

Corrosive. Causes irreversible eye damage. Harmful if swallowed. Do not get in eyes or on clothing. If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical-resistant apron. If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

FIRST AID	
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
<b>HOT LINE NUMBERS</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact CHEMTREC for transportation and spills (800) 424-9300; for Human health (800) 832-HELP (800-832-4357); and ASPCA for animal health (800) 345-4735.	

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Drift or runoff from treated areas may be hazardous to aquatic organisms and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target area. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other spraying. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not

contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.  
**PESTICIDE STORAGE:** Always store pesticides in a secured warehouse or storage building. Do not store near open containers of herbicides and other pesticides, fertilizer and seed. Protect product from freezing. Store at temperatures above 25°F. If allowed to freeze, remix before using. This does not alter the product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.  
**PRODUCT DISPOSAL:** Pesticide wastes are acutely hazardous. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.  
**CONTAINER DISPOSAL FOR NON-REFILLABLE CONTAINERS:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

EPA REG. NO. 228-447-10404  
Distributed by:  
**LESCO, Inc.**, 1301 East 9th Street  
Cleveland, OH 44114-1849

EPA EST. NO. 228-1L-1

LESCO is a registered trademark and the sweeping design is a trademark of LESCO Technologies, LLC. Momentum is a registered trademark and used by LESCO, Inc. with permission of the trademark owner.

PULL HERE TO OPEN  
PRESS TO RESEAL

331190

RV040306